

Deforming Convex Hypersurfaces to a Hypersurface with Prescribed Harmonic Mean Curvature *

Huai-Yu Jian

Department of Applied Mathematics, Tsinghua University, Beijing 100084, P.R.China

(e-mail: hjian@math.tsinghua.edu.cn)

Abstract. Let F be a smooth convex and positive function defined in $A = \{x \in R^{n+1} : R_1 < |x| < R_2\}$ satisfying $F(x) \geq nR_2$ on the sphere $|x| = R_2$ and $F(x) \leq nR_1$ on the sphere $|x| = R_1$. In this paper, a heat flow method is used to deform convex hypersurfaces in A to a hypersurface whose harmonic mean curvature is the given function F .

Keywords: heat flows, prescribed curvature problems, hypersurfaces, convexity-preserving, parabolic equations on manifolds.

1991 AMS Mathematical Subject Classification: 35K, 58G

1. Introduction

Let M be a smooth embedded hypersurface in R^{n+1} and $k_1, k_2, \dots, k_n$ be its principal curvatures. Then H^{-1} is called the harmonic mean curvature of M if

$$H = \frac{1}{k_1} + \frac{1}{k_2} + \dots + \frac{1}{k_n}. \quad (1.1)$$

The question which we are concerned with is that given a function f defined in R^{n+1} , under what conditions does the equation

$$H^{-1}(X) = f(X), X \in M$$

has a solution for a smooth, closed, convex and embedded hypersurface M , where X is a position vector on M .

The kind of such question was proposed by S.T.Yau in his famous problem section [Y]. Many authors have studied the cases of mean curvature and Gauss curvature instead. See, for instance, [BK], [TW], [T], [CNS], and [TS1] for the mean curvature and [O1], [O2], [TS2] for the Gauss curvature, [CNS], [G1] and [G2] for general curvature functions.

*Supported by National Science Foundation of China and by grant of Morningside Center of Mathematics, CAS

Let $F = f^{-1}$, then the problem above is equivalent to looking for a smooth, closed, convex and embedded hypersurface M in R^{n+1} such that

$$H(X) = F(X), X \in M \quad (1.2)$$

where H is the inverse of harmonic mean curvature given by (1.1). We are interested only in the hypersurfaces $M \subset A$, a ring domain defined by

$$A = \{X \in R^{n+1} : R_1 < |X| < R_2\}.$$

for some constants $R_2 > R_1 > 0$. For this purpose, we need to suppose that F is a smooth positive function defined in R^{n+1} satisfying

$$(a) \ F(X) > nR_2 \text{ for } |X| = R_2 \text{ and } F(X) < nR_1 \text{ for } |X| = R_1,$$

and

$$(b) \ F \text{ is concave in } A.$$

We will use a heat flow method to deform convex hypersurfaces to a solution to (1.2). That is we consider the parabolic equation

$$\left. \begin{aligned} \frac{\partial X}{\partial t} &= (H(X) - F(X)) \nu(X), X \in M_t, t \in (0, T) \\ X(\cdot, 0) &\text{ given,} \end{aligned} \right\} \quad (1.3)$$

where $X(x, t) : S^n \rightarrow R^{n+1}$ is the parametrization of M_t given by inverse Gauss map, which will be solved, and $\nu(X)$ is the outer normal at $X \in M_t$, so $\nu(X(x, t)) = x$ by the definition. Of course, M_0 is a given initial hypersurface.

The following is our main result of this paper.

Theorem 1.1. Suppose that F is a smooth positive function satisfying conditions (a) and (b), and a initial hypersurface $M_0 \subset A$ is smooth, uniformly convex and embedded, satisfying $H(X_0) \geq F(X_0)$ for all $X_0 \in M_0$. Then equation (1.3) has a unique smooth solution for $T = \infty$ which parametrizes a family of smooth, closed, uniformly convex and embedded hypersurfaces, $\{M_t : t \in [0, \infty)\}$. Moreover, there exists a subsequence $t_k \rightarrow \infty$ such that M_{t_k} converges to a smooth, closed, uniformly convex and embedded hypersurface which lies in $\bar{A}$ and solves problem (1.2).

We will have to meet two difficulties in proving the result above. One is the gradient estimate for the support function, $u(x, t)$, of the hypersurfaces M_t which solves the equation (2.4) below; the other is the proof of convexity preserving (see the inequality (2.5)). In order to overcome the first difficulty, we use the well-known equality $|X(x, t)|^2 = u^2 + |\nabla u|^2$ and a general geometric result suggested by R. Bartnik to estimate the $|X(x, t)|^2$ for the equation (1.3). The proof of convexity-preserving is inspired by the computations in Hamilton [H1, H2] and Huisken [HU].

Remark 1.2. Applying a usual approximation method, one can replace condition (a) in theorem 1.1 by

$$(a') \ F(X) \geq nR_2 \text{ for } |X| = R_2 \text{ and } F(X) \leq nR_1 \text{ for } |X| = R_1.$$

After completing the paper, the author was aware that similar results had been obtained by Gerhard in [G1]. Although the results in [G1] are very general, they don't

include the above theorem 1.1 because our curvature function H does not belong the class (K) in [G1] and our initial hypersurfaces M_0 may be arbitrary instead of the fixed barrier $M_1 = \{|X| = R_1\}$ in [G1]. Moreover, the arguments are absolutely different.

Acknowledgement. *The author would like to thank professors R. Bartnik, K. Tso and W.Y. Ding for many helpful conversations and professor B. Chow for his interests in this work.*

2. Evolution equations, convexity-preserving and global solutions

In this section, we will at first reduce the equation (1.3) to a equivalent quasilinear parabolic equation on S^n for the support function of M_t , then we will show that this parabolic problem is globally solvable and it preserves convexity.

We recall some facts in [U;p.97-98]. Let $e_1, e_2, \dots, e_n$ be a smooth local orthonormal frame on S^n , and let $\nabla_i = \nabla_{e_i}$, $i=1, 2, \dots, n$ and $\nabla = (\nabla_1, \nabla_2, \dots, \nabla_n)$ be the covariant derivatives and the gradient on S^n , respectively. Since $X(x, t)$ is the inverse Gauss map, the support function of M_t is given by

$$u(x, t) = \langle x, X(x, t) \rangle, x \in S^n \quad (2.1)$$

where $\langle \cdot, \cdot \rangle$ denotes the usual inner product in R^{n+1} . The second fundamental form of M_t is

$$h_{ij}(X(x, t)) = \nabla_i \nabla_j u(x, t) + \delta_{ij} u(x, t), i, j = 1, 2, \dots, n.$$

If M_t is uniformly convex, then h_{ij} is invertible, and hence the inverse harmonic mean curvature is the sum of all the eigenvalues of the matrix

$$b_{ij} = [g^{ik} h_{kj}]^{-1},$$

where g_{ij} is the metric of M_t . But the Gauss-Weingarten relation

$$\nabla_i x = h_{ik} g^{kl} \nabla_l X$$

and the fact $\langle \nabla_i x, \nabla_j x \rangle = \delta_{ij}$ imply $g_{ij} = h_{ik} h_{jk}$. Therefore, $b_{ij} = h_{ij}$ and

$$H = \frac{1}{k_1} + \dots + \frac{1}{k_n} = \Delta u + nu, \quad (2.2)$$

where $\Delta = \sum_{i=1}^n \nabla_i \nabla_i$. Furthermore, since $x, \nabla_1 x, \nabla_2 x, \dots, \nabla_n x$ form a standard orthonormal basis at point $X(x, t)$, so we have

$$\begin{aligned} X(x, t) &= \langle X, x \rangle x + \langle X, \nabla_i x \rangle \nabla_i x \\ &= ux + \nabla_i \langle X, x \rangle \nabla_i x \\ &= ux + \nabla_i u \nabla_i x. \end{aligned} \quad (2.3)$$

Using the results above and repeating the argument [U: p.98-101] in verbatim, one has obtained the following lemma.

Lemma 2.1. If for $t \in [0, T)$ with $T \leq \infty$ $X(x, t)$ is a solution of (1.3) which parametrizes a smooth, closed, uniformly convex and embedded hypersurface, then the support functions $u(x, t)$ of M_t satisfy

$$\left. \begin{aligned} \frac{\partial u}{\partial t} &= \Delta u + nu - F(ux + \nabla_i u \nabla_i x), \quad (x, t) \in S^n \times (0, T) \\ u(\cdot, 0) &= u_0(\cdot), \text{ the support function of } M_0 \end{aligned} \right\} \quad (2.4)$$

and

$$\nabla^2 u + uI > 0 \text{ in } S^n \times (0, T), \quad (2.5)$$

where I denotes the $n \times n$ unit matrix. Conversely, if u is a smooth solution to (2.4) and satisfies (2.5), then the hypersurface M_t , determined by its support function $u(x, t)$, is a smooth, closed, uniformly convex, embedded hypersurface and solves (1.3) for $t \in [0, T)$.

From now on, we assume that the initial hypersurface M_0 is smooth, closed, uniformly convex. That is $u_0 \in C^\infty(S^n)$ and for some positive constant C_0 ,

$$C_0 I \leq \nabla^2 u_0 + u_0 I. \quad (2.6)$$

Noting that (2.4) is a quasilinear parabolic equation on the compact manifold S^n , by standard result for short-time existence (see, for example, [H3]), we have

Lemma 2.2. There exists a maximal existence time $T = T(u_0) \in (0, \infty]$ such that (2.4) has a unique smooth solution $u \in C^\infty(S^n \times (0, T)) \cap C([0, T]; C^\infty(S^n))$. If $T < \infty$, then

$$\lim_{t \rightarrow T^-} \|u(\cdot, t)\|_{C^1(S^n)} = \infty.$$

Remark 2.3. One can use the contraction principle and repeat the same argument as in proving the short-time existence for harmonic heat flows to give a direct proof of this lemma. See [ES] or [D].

For the sake of deriving an apriori estimates, we need the following geometric result which was suggested by R. Bartnik.

Lemma 2.4. Let X be the positive vector of a smooth, closed hypersurface M in R^{n+1} with outer normal $\nu(X)$ at $X \in M$. Then if $|X| = \langle X, X \rangle^{\frac{1}{2}}$ attains a maximum R at a point $X_0 \in M$, then $X_0 = R\nu(X_0)$ and

$$\Pi(w, w) \geq \frac{1}{R} g(w, w), \forall w \in T_{X_0} M;$$

if $|X|$ attains a minimum r at a point $X_0 \in M$, then $X_0 = r\nu(X_0)$ and

$$\Pi(w, w) \leq \frac{1}{r} g(w, w), \forall w \in T_{X_0} M,$$

where g is the metric on M and Π is the second fundamental form of M with respect to the direction $-\nu$.

Proof. We consider only the first case, because the latter is completely analogous. For $X_0 \in M$ and any $w \in T_{X_0} M$, choose a curve $\gamma(s)$ on M , $\gamma : [0, 1] \rightarrow M$, such that

$$\gamma(0) = X_0, \quad \dot{\gamma} = w.$$

Let $\rho(X) = |X|$. since $\rho^2(X)$ attains a maximum at $X_0 \in M$, then at this point

$$\nabla \rho^2 = 0 \quad \text{and} \quad \nabla^2 \rho^2 \leq 0.$$

Therefore, we have

$$\frac{d}{ds} \rho^2(\gamma(0)) = \nabla \rho^2(X_0) \cdot \dot{\gamma}(0) = 0$$

and

$$\frac{d^2}{ds^2} \rho^2(\gamma(0)) = \dot{\gamma}(0) \cdot \nabla^2 \rho^2(X_0) \cdot \dot{\gamma}(0) + \nabla \rho^2(X_0) \cdot \ddot{\gamma}(0) \leq 0.$$

On the other hand,

$$\frac{d\rho^2}{ds} = \frac{d}{ds} |\gamma(s)|^2 = 2\gamma(s) \cdot \dot{\gamma}(s)$$

and

$$\frac{1}{2} \frac{d^2 \rho^2}{ds^2} = |\dot{\gamma}(s)|^2 + \gamma(s) \cdot \ddot{\gamma}(s).$$

Thus,

$$0 = 2\gamma(0) \cdot \dot{\gamma}(0) = 2X_0 \cdot w \tag{2.7}$$

and

$$0 \geq |\dot{\gamma}(0)|^2 + X_0 \cdot \ddot{\gamma}(0). \tag{2.8}$$

Since $w \in T_{X_0}M$ can be arbitrary, (2.7) implies

$$X_0 = |X_0| \nu(X_0) = R\nu(X_0),$$

and so,

$$\begin{aligned} X_0 \cdot \ddot{\gamma}(0) &= R \langle \nu(X_0), \ddot{\gamma}(0) \rangle \\ &= R \langle \nu(X_0), D_{\dot{\gamma}(0)} \dot{\gamma}(0) \rangle \\ &= -R \Pi(\dot{\gamma}(0), \dot{\gamma}(0)) \\ &= -R \Pi(w, w), \end{aligned}$$

which, together with (2.8), gives us that

$$\begin{aligned} \Pi(w, w) &\geq \frac{1}{R} |\dot{\gamma}(0)|^2 \\ &= \frac{1}{R} \langle \dot{\gamma}(0), \dot{\gamma}(0) \rangle \\ &= \frac{1}{R} \langle D_{\dot{\gamma}(0)} X, D_{\dot{\gamma}(0)} X \rangle \\ &= \frac{1}{R} g(\dot{\gamma}(0), \dot{\gamma}(0)) \\ &= \frac{1}{R} g(w, w). \end{aligned}$$

Lemma 2.5. Suppose that $M_0 \subset\subset A$ in addition to (2.6). Let u be a smooth solution to (2.4) and satisfies (2.5) on $S^n \times (0, T)$ with $T \leq \infty$. Then for all $(x, t) \in S^n \times (0, T)$, we have

$$R_1^2 < u^2(x, t) + |\nabla u(x, t)|^2 < R_2^2.$$

Proof. It follows from lemma 2.1 that the position vector $X(\cdot, t)$ of M_t determined by the support function $u(x, t)$ satisfies (1.3), i.e.

$$\left. \begin{aligned} \frac{\partial X}{\partial t} &= (H(X) - F(X))\nu(X), X(x, t) \in M_t, (x, t) \in S^n \times (0, T) \\ X(\cdot, 0) &= X_0. \end{aligned} \right\} \quad (2.9)$$

Moreover, (2.3) and the fact that $x, \nabla_1 x, \dots, \nabla_n x$ form a standard orthonormal basis imply that

$$|X|^2 = u^2 + |\nabla u|^2, \forall (x, t) \in S^n \times (0, T). \quad (2.10)$$

Thus, it is sufficient to prove that for X solving (2.9),

$$R_1^2 < |X(x, t)|^2 < R_2^2, \forall (x, t) \in S^n \times (0, T). \quad (2.11)$$

For each $t \in [0, T)$, let

$$P_{min}(t) = \min_{X \in M_t} |X|^2 = \min_{x \in S^n} |X(x, t)|^2,$$

and

$$P_{max}(t) = \max_{X \in M_t} |X|^2 = \max_{x \in S^n} |X(x, t)|^2.$$

By virtue of the assumption for M_0 , we have (2.6) and

$$R_1^2 < P_{min}(0) \leq P_{max}(0) < R_2^2. \quad (2.12)$$

Since M_t is smooth, P_{min} and P_{max} are obviously Lipschitzian on $[0, T)$. Were the inequality (2.11) not true, then by (2.12) we could find t_1 and t_2 in $[0, T)$ such that either

$$P_{min}(t_1) = R_1^2 \quad (2.13)$$

or

$$P_{max}(t_2) = R_2^2. \quad (2.14)$$

Without loss of generality, we assume that the case (2.13) happens, and the case (2.14) is completely similar. Let

$$t^* = \inf\{t \in (0, T) : P_{min}(t) = R_1^2\},$$

and choose $x^* \in S^n$ such that

$$P_{min}(t^*) = |X(x^*, t^*)|^2.$$

In order to compute the principal curvatures $k_1, \dots, k_n$ of M_{t^*} at $X(x^*, t^*)$, we use the principal direction $\xi_1, \dots, \xi_n$ to obtain

$$\Pi(\xi_i, \xi_i) = k_i \langle \xi_i, \xi_i \rangle = k_i g(\xi_i, \xi_i),$$

so

$$k_i = \Pi(\xi_i, \xi_i)g^{-1}(\xi_i, \xi_i), i = 1, 2, \dots, n.$$

Thus, lemma 2.4 implies that $X(x^*, t^*) = R_1 \nu(X(x^*, t^*))$ and

$$k_i \leq \frac{1}{R_1}, \quad i = 1, 2, \dots, n$$

at $X(x^*, t^*)$. Therefore, we have

$$H(X(x^*, t^*)) \geq nR_1,$$

and

$$\begin{aligned} \frac{\partial |X|^2}{\partial t}(x^*, t^*) &= X \cdot \frac{\partial X}{\partial t} \\ &= R_1 \nu \cdot (H - F) \nu \\ &\geq R_1 [nR_1 - F(X(x^*, t^*))] \\ &> 0, \end{aligned}$$

where we have used the condition (a) for $X(x^*, t^*) = R_1$.

On the other hand, we obviously have

$$|X(x^*, t)|^2 > |X(x^*, t^*)|^2, \forall t \in [0, t^*)$$

thus

$$\frac{\partial |X|^2}{\partial t}(x^*, t^*) \leq 0$$

which yields a contradiction.

Next, we will prove that the convexity of M_t is preserved. That is, (2.5) remains true for all $t \in (0, T)$ if it is so at $t = 0$.

Let

$$h_{kl} = \nabla_k \nabla_l u + \delta_{kl} u, \quad k, l = 1, 2, \dots, n.$$

It follows directly from (2.4) and (2.3) that

$$\frac{\partial}{\partial t}(\delta_{kl} u) = \delta_{kl} \left[\sum_{i=1}^n h_{ii} - F(ux + X(x, t)) \right]. \quad (2.15)$$

On the other hand, we differentiate the equation (2.4) twice to get

$$\frac{\partial}{\partial t} \nabla_k \nabla_l u = \nabla_k \nabla_l \Delta u + n \nabla_k \nabla_l u - F_j(X) \nabla_k X^j - F_{jh}(X) \nabla_k X^h \nabla_l X^j. \quad (2.16)$$

Using the standard formula for interchanging the order of covariant differentiation with respect to the orthonormal frame on S^n , we have

$$\nabla_k \nabla_l \Delta u = \Delta \nabla_k \nabla_l u - 2n \nabla_k \nabla_l u + 2\delta_{kl} \Delta u, \quad (2.17)$$

see, for instance, [CLT, p.85]. Thus, combining (2.15)-(2.17), we get

$$\begin{aligned} \frac{\partial}{\partial t} h_{kl} &= \Delta h_{kl} + \delta_{kl} \Delta u - n \nabla_k \nabla_l u + \delta_{kl} \left[\sum_{i=1}^n h_{ii} - F(X(x, t)) \right] \\ &\quad - F_j(X(x, t)) \nabla_k \nabla_l X^j - F_{jh}(X) \nabla_k X^h \nabla_l X^j. \end{aligned} \quad (2.18)$$

Set

$$G(x, t) = \sum_{i=1}^n h_{ii}(x, t) - F(X(x, t)).$$

By virtue of (2.18), (2.3), and (2.4), we see that

$$\begin{aligned} \frac{\partial G}{\partial t} &= \frac{\partial}{\partial t} \sum_{i=1}^n h_{ii} - \frac{\partial F}{\partial t} \\ &= \Delta G + nG - \frac{\partial F}{\partial t} \\ &= \Delta G + nG - F_j(X) \left(\frac{\partial u}{\partial t} x^j + \nabla_i \frac{\partial u}{\partial t} \nabla_i x^j \right) \\ &= \Delta G + nG - F_j(X) (Gx^j + \nabla_i G \nabla_i x^j). \end{aligned} \quad (2.19)$$

Lemma 2.6. Suppose that in addition to (2.6), the support function u_0 of the initial hypersurface M_0 satisfies

$$G(x, 0) = \Delta u_0(x) + nu_0(x) - F(u_0 + \nabla_i u_0 \nabla_i x) \geq 0, \quad x \in S^n \quad (2.20)$$

and let $u(x, t)$ be a smooth solution to (2.4) on $S^n \times (0, T)$ with $T \leq \infty$. Then for all $(x, t) \in S^n \times (0, T)$,

$$G(x, t) = \Delta u(x, t) + nu(x, t) - F(ux + \nabla_i u \nabla_i x) \geq 0. \quad (2.21)$$

Proof. It is sufficient to prove that for each $T_0 < T$, (2.21) holds true for all $(x, t) \in S^n \times [0, T_0)$. Let

$$C_1 = \max \{ |u(x, t)| + |\nabla u(x, t)| : (x, t) \in S^n \times [0, T_0] \},$$

$$C_2 = \max \left\{ \sum_{j=1}^n \left| \frac{\partial F}{\partial y_j} \right| : |y| \leq C_1 \right\},$$

and

$$G_{\min}(t) = \min \{ G(x, t) : x \in S^n \}.$$

If $G_{\min}(t) \leq 0$ for some $t \in (0, T_0)$ we could find $x_t \in S^n$ such that

$$G(x_t, t) = G_{\min}(t) \leq 0, \quad \Delta G(x_t, t) \geq 0, \quad \nabla G(x_t, t) = 0.$$

Thus, (2.19) implies that at (x_t, t)

$$\frac{\partial G}{\partial t} \geq nG - F_j(X) Gx^j \geq (n + C_2)G.$$

This yields

$$\frac{\partial}{\partial t} \left(G e^{-(n+C_2)t} \right) (x_t, t) \geq 0.$$

Note that $e^{-(n+C_2)t} G_{\min}(t)$ is a Lipchitzian function of t . Denote it by $B(t)$ for simplicity. Then the result above gives us that

$$\liminf_{\tau \rightarrow 0^+} \frac{B(t + \tau) - B(t)}{\tau} \geq 0 \quad \text{when} \quad B(t) \leq 0 \quad \text{for} \quad t \in (0, T_0).$$

Also see [U, p.107]. Now by a result of Hamilton[H2, lemma 3.1] we conclude that

$$B(t) \geq 0, \quad \forall t \in [0, T_0].$$

This proves the lemma.

Lemma 2.7. Assume that the support function u_0 of the initial hypersurface M_0 satisfies (2.6) and (2.20). Let $u(x, t)$ be a smooth solution to (2.4) on $S^n \times (0, T)$ with $T \leq \infty$. Then for all $(x, t) \in S^n \times (0, T)$, we have

$$\nabla^2 u(x, t) + u(x, t)I > 0.$$

Proof. If the conclusion were not true, we could find a finite number $t_0 \in (0, T)$ such that the minimum eigenvalue of the matrix $[h_{kl}(x, t_0)]$ is zero, but $[h_{kl}(x, t)]$ is positive definite for all $(x, t) \in S^n \times [0, t_0]$. Thus the inverse matrix $[h^{pq}(x, t)]$ exists for all such (x, t) . Let

$$g_{kl}(x, t) = \langle \nabla_k X(x, t), \nabla_l X(x, t) \rangle, \quad k, l = 1, 2, \dots, n$$

be the metrix of M_t . Then Gauss-weingarten relation and the fact $\delta_{kl} = \langle \nabla_i x, \nabla_j x \rangle$ gives us that

$$\nabla_i x^j = h_{ik} g^{kl} \nabla_l X^j, \quad g_{ij} = h_{ik} h_{jk}.$$

See [U, p.98]. Therefore, for each $(x, t) \in S^n \times [0, t_0]$, we have

$$\begin{aligned} \nabla_k \nabla_l X^j &= \nabla_k (g_{lm} h^{mi} \nabla_i x^j) \\ &= \nabla_k (h_{li} \nabla_i x^j) \\ &= \nabla_k \nabla_i x^j h_{il} + \nabla_i x^j \nabla_k h_{li}. \end{aligned}$$

On the other hand, it easily follows from the standard formula for commuting the order of covariant differentiation on S^n that

$$\nabla_i h_{jk} = \nabla_j h_{ik} = \nabla_k h_{ij} \quad \text{for all } i, j = 1, 2, \dots, n$$

see [U, p.111]. With the aid of two equatities above, (2.18) turns to

$$\begin{aligned} \frac{\partial}{\partial t} h_{kl} &= \Delta h_{kl} + \delta_{kl} \Delta u - n \nabla_k \nabla_l u + \delta_{kl} G \\ &\quad - F_j(X) (\nabla_k \nabla_i x^j h_{il} + \nabla_i x^j \nabla_i h_{lk}) - F_{jh}(X) \nabla_k X^h \nabla_l X^j. \end{aligned} \quad (2.22)$$

Now let us suppose that the minimum eigenvalue of $[h_{kl}]$ over S^n at time $t \in [0, t_0]$ attains at a point $x_t \in S^n$. By rotating the frame $e_1, e_2, \dots, e_n$, we may assume that $h_{11}(x_t, t)$ is the minimum eigenvalue and $h_{j1}(x_t, t) = 0$ for $j = 2, 3, \dots, n$. (Also see [CLT, p.89].) Combining (2.22) and lemma 2.6 together, at (x_t, t) we have

$$\frac{\partial h_{11}}{\partial t} \geq \Delta u - n \nabla_1 \nabla_1 u - F_j(X) \nabla_1 \nabla_1 x^j h_{11} - F_{jh} \nabla_1 X^j \nabla_1 X^h, \quad (2.23)$$

where we have used $\Delta h_{11} \geq 0$ and $\nabla h_{11} = 0$ at (x_t, t) . Since at (x_t, t)

$$nu + n\nabla_1 \nabla_1 u \leq \sum_{i=1}^n h_{ii} = \Delta u + nu,$$

we have

$$\Delta u - n\nabla_1 \nabla_1 u \geq 0.$$

Applying this equality and the concavity of F to (2.23), we obtain

$$\frac{\partial h_{11}}{\partial t} \geq -F_j(X) \nabla_1 \nabla_1 x^j h_{11}.$$

Denote

$$C_3 = \max \{|u(x, t)| + |\nabla u(x, t)| : (x, t) \in S^n \times [0, t_0]\},$$

$$C_4 = \max \left\{ \sum_{j=1}^n |F_j(Y)| : |Y| \leq C_3 \right\},$$

and

$$C_5 = C_4 \max \{|\nabla_1 \nabla_1 x| : x \in S^n\}.$$

Noting $h_{11}(x_t, t)$ is positive for all $t \in [0, t_0)$, we have

$$\frac{\partial}{\partial t} h_{11}(x_t, t) \geq -C_5 h_{11}(x_t, t), \quad t \in [0, t_0).$$

That is

$$\frac{\partial}{\partial t} (h_{11}(x_t, t) e^{C_5 t}) \geq 0, \quad t \in [0, t_0).$$

By the maximum principle and the condition (2.6), we see that

$$h_{11}(x_t, t) \geq h_{11}(x_0, 0) e^{-C_5 t} \geq C_0 e^{-C_5 t} \quad \forall t \in [0, t_0).$$

Letting $t \rightarrow t_0^-$, we obtain

$$h_{11}(x_{t_0}, t_0) \geq C_0 e^{-C_5 t_0},$$

which is contradictory with the assumption that the minimum eigenvalue of $[h_{kl}]$ is zero. This proves the lemma

Theorem 2.8. Suppose that the initial hypersurface $M_0 \subset\subset A$ and its support function u_0 satisfies (2.6) and (2.20). Then there exists a unique smooth solution u to the following problem:

$$\left. \begin{aligned} \frac{\partial u}{\partial t} &= \Delta u + nu - F(ux + \nabla_i u \nabla_i x) \quad \text{in } S^n \times (0, \infty) \\ u(\cdot, 0) &= u_0(\cdot) \quad \text{on } S^n, \end{aligned} \right\} \quad (2.24)$$

Moreover, for all $(x, t) \in S^n \times (0, \infty)$, $u(x, t)$ satisfies

$$\nabla^2 u + uI > 0, \quad (2.25)$$

$$\frac{\partial u}{\partial t} = \Delta u + nu - F(ux + \nabla_i u \nabla_i x) \geq 0, \quad (2.26)$$

and

$$R_1^2 < u^2 + |\nabla u|^2 < R_2^2. \quad (2.27)$$

Proof. By virtue of lemma 2.2, we know that the problem (2.24) has a unique smooth solution on $S^n \times (0, T)$ with some $T \leq \infty$. Moreover, lemma 2.6 and 2.7 tell us that both (2.25) and (2.26) are satisfied on $S^n \times (0, T)$. Thus, by lemma 2.5, we see that (2.27) is true for all $(x, t) \in S^n \times (0, T)$. Using lemma 2.2 again, we know that T is nothing but ∞ . This completes the theorem.

3. Converging to a convex hypersurface

In this section, we will use the theorem 2.8 in last section to prove the main result of this paper, theorem 1.1.

We begin with choosing a smooth, closed, uniformly convex hypersurface M_0 such that its support function satisfies (2.6) and (2.20). (The existence of such M_0 is obvious due to the condition (a)). By theorem 2.8 and lemma 2.1 we obtain a family of smooth, closed, uniformly convex hypersurfaces M_t whose position vectors are

$$X(x, t) = u(x, t)x + \nabla_i u(x, t)\nabla_i x, \quad (x, t) \in S^n \times [0, \infty),$$

where $u(x, t)$ are the support functions of M_t and satisfy (2.24)-(2.27) in theorem 2.8. It follows from (2.26) and (2.24) that

$$\left(\frac{\partial u}{\partial t}\right)^2 = \frac{\partial u}{\partial t}(\Delta u + nu - F(X(x, t))) \leq \frac{\partial u}{\partial t}(\Delta u + nu).$$

Hence

$$\int_{S^n} \left(\frac{\partial u}{\partial t}\right)^2 dx \leq \frac{1}{2} \frac{\partial}{\partial t} \int_{S^n} (nu^2 - |\nabla u|^2) dx, \quad \forall t \in (0, \infty).$$

This, together with (2.27), implies that

$$\int_0^\infty \int_{S^n} \left(\frac{\partial u}{\partial t}\right)^2 dx dt \leq (n+2)R_2^2 \cdot \text{vol}(S^n). \quad (3.1)$$

Since (2.27) implies that for each $\alpha \in (0, 1)$

$$nu - F(ux + \nabla_i u \nabla_i x) \in L^\infty(S^n \times [0, \infty)),$$

it follows from a property of heat equation that

$$\|u(\cdot, t)\|_{C^{1,\alpha}(S^n)} \leq C \quad (3.2)$$

uniformly in $t \in [1, \infty)$. See [LSU, ch.4] or [D]. Moreover, by (3.2) and the same argument, we have

$$\|u(\cdot, t)\|_{C^{3,\alpha}(S^n)} \leq C \quad (3.3)$$

uniformly in $t \in [1, \infty)$. Using (3.1) and (3.3), we can find a sequence $t_k \rightarrow \infty$ as $k \rightarrow \infty$ such that

$$\int_{S^n} \left(\frac{\partial u}{\partial t_k}\right)^2 dx \rightarrow 0, \quad (3.4)$$

and

$$u(\cdot, t_k) \rightarrow U_0 \text{ in } C^{3,\alpha}(S^n). \quad (3.5)$$

Furthermore, (2.24), (3.4) and (3.5) gives us

$$\int_{S^n} [\Delta U_0 + nU_0 - F(U_0x + \nabla_i U_0 \nabla_i x)]^2 dx = 0,$$

which yields

$$\Delta U_0 + nU_0 - F(U_0x + \nabla_i U_0 \nabla_i x) = 0, \quad x \in S^n. \quad (3.6)$$

Since $U_0 \in C^2(S^n)$, applying a standard elliptic theory to the equation (3.6), we know $U_0 \in C^\infty(S^n)$.

On the other hand, it easily follows from (3.4), (3.5) and (3.6) that as $k \rightarrow \infty$

$$\frac{\partial u}{\partial t_k}(\cdot, t_k) \rightarrow 0 \text{ in } C(S^n).$$

Hence

$$\nabla \frac{\partial u}{\partial t_k}(\cdot, t_k) \rightarrow 0 \text{ in } C(S^n). \quad (3.7)$$

Therefore, (3.5) and (3.7) implies that

$$X(x, t_k) = u(x, t_k)x + \nabla_i u(x, t_k) \nabla_i x \rightarrow U_0x + \nabla_i U_0 \nabla_i x \text{ in } C^1(S^n), \quad (3.8)$$

and

$$\frac{\partial X}{\partial t_k}(x, t_k) = \frac{\partial u}{\partial t_k}(x, t_k)x + \nabla_i \frac{\partial u}{\partial t_k}(x, t_k) \nabla_i x \rightarrow 0 \text{ in } C(S^n). \quad (3.9)$$

Since each hypersurface M_t is smooth, closed, uniformly convex, lemma 2.1 says that $X(x, t)$ satisfy (1.3) for all $t \in [0, \infty)$. Thus, using (2.2), (3.5), (3.8) and (3.9), we know that

$$Y_0(x) = U_0(x)x + \nabla_i U_0 \nabla_i x \quad x \in S^n \quad (3.10)$$

satisfies the equation (1.2).

Now let M_y be the hypersurface determined by the position vector $Y_0(x)$, $x \in S^n$. Obviously, M_y is closed and encloses the origin, and is convex because of (3.5), (3.8) and the uniform convexity of hypersurfaces M_t for any $t \in [0, \infty)$. So M_y is uniformly convex because Y_0 satisfies (1.2). Furthermore, due to the fact $U_0 \in C^\infty(S^n)$, M_y is smooth, too. Finally, using (2.27), (3.5) and (3.10), we obtain

$$R_1^2 \leq |X_0(x)|^2 \leq R_2^2, \quad x \in S^n,$$

which means that the hypersurface M_y lies in $\bar{A}$. This prove theorem 1.1.

References

- [BK] L. Bakelman and B. Kantor, Existence of spherically homeomorphic hypersurfaces in Euclidean space with prescribed mean curvature, *Geometry and Topology*, Leningrad(1974), 3-10.
- [CNS] L. Caffarelli, J. Nirenberg and J. Spruck, The Dirichlet problem for nonlinear second order elliptic equation. IV: Starshaped compact Weingarten hypersurfaces, *Current topics in partial differential equations*, ED. by Y. Ohya, K. Kasahara, N. Shimakura (1986), 1-26, Kinokuniya Co., Tokyo.
- [CLT] B. Chow, L.P. Liou and D.H. Tsai, On the nonlinear parabolic equation $\partial_t u = F(\Delta u + nu)$ on S^n , *Comm. Anal. Geom.* 3(4)(1996), 75-94.
- [D] W.Y. Ding, *Lecture Notes on harmonic maps*, Inst. Math. Academia Sinica, 1995.
- [G1] C. Gerhardt, Closed weingarten hypersurfaces in Riemannian manifolds, *J. Differ. Geom.* 43(1996), 612-641.
- [G2] C. Gerhardt, Hypersurfaces of prescribed weingarten curvature, *Math. Z.* 224(1997), 167-194.
- [ES] J. Eells and J.H. Sampson, Harmonic mappings of Riemannian manifolds, *Amer. J. Math.* 86(1964), 109-160.
- [H1] R. S. Hamilton, Three-manifolds with positive Ricci curvature, *J. Differ. Geom.* 17(1982), 255-306.
- [H2] R. S. Hamilton, Four-manifolds with positive Ricci curvature, *J. Differ. Geom.* 24(1986), 153-179.
- [H3] R.S. Hamilton, *Lecture Notes on heat equation in geometry*, Honolulu, Hawaii, 1989.
- [HU] G. Huisken, Flow by mean curvature of convex hypersurfaces into spheres, *J. Differ. Geom.* 20(1984), 237-266.
- [LSU] O. Ladyzhenskaja, A. Solonnikov and N. N. Uralceva, *Linear and quasilinear equations of parabolic type*, Trans. Amer. Math. Soc. 23, Providence, 1968.
- [O1] V.I. Oliker, Hypersurfaces in R^{n+1} with prescribed Gaussian curvature and related equations of Morge-Ampere type, *Comm. p.d.e.*, 9(1984), 807-839.
- [O2] V.I. Oliker, The problem of embedding S^n into R^{n+1} with prescribed Gauss curvature and its solution by variational methods, *Trans. Amer. Math. Soc.*, 295(1986), 291-303.
- [T] A. Treibergs, Existence and convexity of hypersurfaces of prescribed curvature, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* 12(1985), 225-241.
- [TW] A. Treibergs and S.W. Wei, Embedded hypersurfaces with prescribed mean curvature, *J. Differ. Geom.* 18(1983), 513-521.
- [TS1] K. Tso, On the existence of convex hypersurfaces with prescribed mean curvature, *Ann. Scuola Norm. Sup. Pisa* 16 (1989), 225-243.
- [TS2] K. Tso, Existence of convex hypersurfaces with prescribed Gauss-Kronecker curvature, *J. Differ. Geom.* (1991), 389-410.
- [U] J.I.E. Urbas, An expansion of convex hypersurfaces, *J. Differ. Geom.* 33(1991), 91-125; Correction to, *ibid.* 35, 763-765.
- [Y] S.T. Yau, Problem section, *Seminar on differential geometry*, Ed. S.T. Yau, *Ann. of Math. Stud.*, Princeton Univ. Press. 102, 1982, 669-706.